

RESEARCH ARTICLE

Open Access

BMP-2 functions independently of SHH signaling and triggers cell condensation and apoptosis in regenerating axolotl limbs

Jean-Charles Guimond², Mathieu Lévesque², Pierre-Luc Michaud¹, Jérémie Berdugo¹, Kenneth Finnsen³, Anie Philip³, Stéphane Roy^{1,2*}

Abstract

Background: Axolotls have the unique ability, among vertebrates, to perfectly regenerate complex body parts, such as limbs, after amputation. In addition, axolotls pattern developing and regenerating autopods from the anterior to posterior axis instead of posterior to anterior like all tetrapods studied to date. Sonic hedgehog is important in establishing this anterior-posterior axis of limbs in all tetrapods including axolotls. Interestingly, its expression is conserved (to the posterior side of limb buds and blastemas) in axolotl limbs as in other tetrapods. It has been suggested that *BMP-2* may be the secondary mediator of sonic hedgehog, although there is mounting evidence to the contrary in mice. Since *BMP-2* expression is on the anterior portion of developing and regenerating limbs prior to digit patterning, opposite to the expression of sonic hedgehog, we examined whether *BMP-2* expression was dependent on sonic hedgehog signaling and whether it affects patterning of the autopod during regeneration.

Results: The expression of *BMP-2* and *SOX-9* in developing and regenerating axolotl limbs corresponded to the first digits forming in the anterior portion of the autopods. The inhibition of sonic hedgehog signaling with cyclopamine caused hypomorphic limbs (during development and regeneration) but did not affect the expression of *BMP-2* and *SOX-9*. Overexpression of *BMP-2* in regenerating limbs caused a loss of digits. Overexpression of *Noggin* (*BMP* inhibitor) in regenerating limbs also resulted in a loss of digits. Histological analysis indicated that the loss due to *BMP-2* overexpression was the result of increased cell condensation and apoptosis while the loss caused by *Noggin* was due to a decrease in cell division.

Conclusion: The expression of *BMP-2* and its target *SOX-9* was independent of sonic hedgehog signaling in developing and regenerating limbs. Their expression correlated with chondrogenesis and the appearance of skeletal elements has described in other tetrapods. Overexpression of *BMP-2* did not cause the formation of extra digits, which is consistent with the hypothesis that it is not the secondary signal of sonic hedgehog. However, it did cause the formation of hypomorphic limbs as a result of increased cellular condensation and apoptosis. Taken together, these results suggest that *BMP-2* does not have a direct role in patterning regenerating limbs but may be important to trigger condensation prior to ossification and to mediate apoptosis.

Background

Bone morphogenetic proteins (*BMPs*) are members of the transforming growth factor- β superfamily and were first discovered when ectopic cartilage and bone formation were induced by demineralized bone matrix implanted into soft tissues of animals [1,2]. However,

their biological roles go far beyond osteogenesis [3,4]. *BMPs* have been extensively studied during the development of the vertebrate limb, which has long been recognized as an excellent experimental system to study the genes and signaling pathways involved in patterning complex structures.

The first stage in which *BMPs* were assessed in limb development is the establishment of the anterior-posterior limb axis. In amniotes, the posterior region of the forming limb bud is referred to as the zone of

* Correspondence: stephane.roy@umontreal.ca

¹Department of Stomatology, Faculty of Dentistry, Université de Montréal, Montreal (QC) Canada

polarizing activity (ZPA) and is an organizer of the anterior-posterior patterning of developing limbs [5]. When cells from the ZPA are grafted into the anterior region of a chick limb bud, polydactyly is induced [6,7]. Previous studies have shown that sonic hedgehog (*Shh*) is expressed exclusively in the ZPA and that implanting *Shh* expressing cells, or beads of *Shh*, in the anterior limb bud is sufficient to reproduce the polydactyly induced with ZPA transplants [5,8]. On the other hand, knockout mice for *Shh* display a major deficiency in the anterior-posterior patterning of their autopods which resemble a spike like structure [9]. Thus, *Shh* appears to be a mediator for the patterning of the anterior-posterior axis of the limb [5,8,10]. Because of the restricted expression of *Shh* to the posterior margin, it remains unclear, however, whether *Shh* induces the limb polarity directly or through a secondary signal. *BMP-2* has received much attention as a downstream target of *Shh*. Indeed, *BMP-2* expression overlaps spatiotemporally with the expression of *Shh* in the early developing limb bud of mammalian and avian embryos [11-15] and can be induced in chick anterior wing buds by ectopic *Shh* expression [12,13]. However, overexpression of *BMP-2* alone does not reproduce the polydactyly obtained with *Shh* overexpression in chick embryos [16,17]. Moreover, *BMP-2* expression is maintained in the limb buds of *Shh* knockout mice [9]. In addition, the polarity and digit identity is normal in mice with *BMP-2* deficient limbs [18]. Hence, these data show that *BMP-2* is not essential to relay *Shh* signaling in developing limbs.

In more advanced limb developmental stages, *BMP-2*, *BMP-4* and *BMP-7* are all expressed in the interdigital (ID) tissue [11,15,19]. Extirpation experiments of ID tissues have shown that anterior-posterior identity is not a fixed property of digital primordia and that the ID tissues are important for digit patterning [20]. Experiments with inhibitors of *BMP* signaling in ID tissue suggested that differential levels of *BMPs* in the ID mesenchyme may be important in directing digit patterning [20]. However, a recent study [18] with conditional knockout for *BMP-2*, *BMP-4* or *BMP-7* showed that these *BMPs* are not essential as signals from the ID mesenchyme for specifying digit identity but that they mediate ID programmed cell death as previously reported [21,22]. In the chicken, overexpression of dominant negative type I-A or type I-B *BMP* receptors (*dnBMPR-IA* or *dnBMPR-IB*), in embryonic hind limbs, reduced ID tissue apoptosis and resulted in webbed feet [23,24]. On the contrary, application of *BMP-2* or *BMP-7* in the ID tissues accelerated and intensified apoptosis [25]. In ducks [19] and bats [26], *BMPs* are also expressed in the ID tissues of developing digits. However, these animals have webbed autopods and apoptosis is absent in

ID tissues, suggesting expression of *BMPs* could have other roles in digit formation.

One aspect that is clearly affected by the expression of *BMP-2* in developing limbs is skeletal development [16,17,27-29]. It was shown, in chicks, that *BMPs* are needed for chondroprogenitor cell determination and/or condensation and subsequent differentiation into chondrocytes [30]. In mice with *BMP-2/BMP-4* deficient limbs, differentiation of condensed prechondrocytes into chondrocytes is delayed, indicating that a threshold level of *BMPs* is required to trigger chondrogenesis [18]. In these mutants, the two posterior digits fail to form as a consequence of the misexpression of the *SOX-9* gene. *SOX-9*, a High Mobility Group (HMG)-domain transcription factor, is the earliest marker of chondrogenesis [31], and a regulator of the type II collagen gene, which is a major constituent of cartilage [29,32]. *SOX-9* is a known target of *BMP-2* [33] and overexpression of *BMP-2* in developing chick limbs induces *SOX-9* expression which then stimulates cartilage formation [29].

Among tetrapod vertebrates, urodele amphibians (i.e. axolotls) have a unique developmental pattern of their limbs. Indeed, while amniotes and anurans develop their digits almost simultaneously with a slightly accelerated onset of posterior digit formation (digit 4 being the first digit to condense in mice) [34-36], urodeles show a reversed sequence with a more progressive/successive development of anterior digits followed by posterior digits [36,37]. Although the order of digit development is different between the aforementioned animals, the end point is the same with similar digit identity in the mature limb [38]. Moreover, even though there is no clearly defined ZPA *per se* in axolotl and newt, *Shh* is expressed, as in amniotes, in the posterior mesenchyme of their developing and regenerating limb buds [39-41]. The inhibition of *Shh* signaling with cyclopamine leads to an anterior-posterior truncation of autopod elements in the developing and regenerating axolotl limbs [42,43]. In addition, overexpression of *Shh* in the anterior mesenchyme of regenerating limbs induces polydactyly [44], showing that as in amniotes, *Shh* has a morphogenetic property in patterning the anterior-posterior axis in developing and regenerating urodele limbs.

The present study addresses the role of *BMP-2* during limb regeneration in axolotl. We report the first cloning of an axolotl full length *BMP* cDNA (*BMP-2*). We use cyclopamine, a pharmacological inhibitor of the hedgehog signaling pathway, to show that, as in mouse, *BMP-2* is not likely to be the secondary signal of *Shh*. *BMP-2* is expressed in the ID tissues of developing digits in virtually all tetrapod species [11,15,19,24,26,45,46] and the expression of *SOX-9* in these digits is linked to *BMP-2* expression [29,30,47,48]. The fact that the sequential development/regeneration of digits in axolotl limbs is

the opposite of that observed in other tetrapods (anterior digits form first in axolotls instead of the posterior digits), even though *Shh* expression is expressed on the posterior side of axolotl developing/regenerating limbs, allowed us to assess whether the expression of *BMP-2* and *SOX-9* correlated with digits appearance (as they condense) or whether their expression correlated with *Shh* signaling in conditions where it was normal or inhibited. The effect of *BMP-2* or *Noggin* (a general *BMP* inhibitor) overexpression on regenerating digits was also assessed.

Results

Cloning and analysis of *BMP-2*

The axolotl full-length cDNA encoding *BMP-2* was cloned, sequenced and compared to the sequence of other vertebrates. Several *BMP-2* clones were isolated from an axolotl cDNA library. The longest clone of 2228 bp [GenBank: EU339232.1] comprising an open reading frame (ORF) of 1188 bp was selected. The analysis of this ORF, using NCBI-BLAST, showed a highly conserved sequence homology of the *BMP-2* ORF with that of other species: 83% identity to human; 83% to mouse; and 67.3% to *Xenopus laevis*. Moreover, the 3' UTR of the *BMP-2* gene was highly conserved with that of other species (data not shown).

The translated ORF of axolotl *BMP-2* has a predicted amino acid sequence of 395 amino acids (Additional File 1: Figure S1) that varies by less than 1% in length when compared to that of human, mouse and *Xenopus laevis*. The amino acid sequence derived from the ORF also showed a highly conserved protein homology between species: 73% with human, 72% with mouse, and 68% with *Xenopus laevis* and resulted in an expected value (E-value) of less than $1e^{-164}$ when blasted in Gen Bank. Seven highly conserved cysteine residues characteristic to all members of the *TGF- β* super family [49,50] are also present (Additional File 1: Figure S1 shaded amino acids).

BMP-2 and *SOX-9* expression during limb development

Whole-mount *in situ* hybridization was performed on developing forelimbs of axolotl from developmental stages 41 to 53. The nomenclature of Nye et al. [51] was used to determine the staging and to label the digits of the forelimb. Digit 1 is the anterior most (corresponding to the index side in human) and digit 4 the posterior most.

At stage 47, the expression of *BMP-2* was mainly detected as a narrow band that surrounded the anterior border of the limb bud and extended slightly to the posterior part of the distal end. Prior to stage 47, the expression of *BMP-2* was expressed as a band at the edge of the limb covering both the anterior to posterior

portions of the limb (data not shown) as described previously for *Xenopus laevis* [52]. *SOX-9* was also expressed in the anterior portion of the limb bud at stage 47 while *Shh* expression is restricted to the posterior (i.e. opposite to that of *BMP-2*; Figure 1). At stage 48, the expression of *BMP-2* had become restricted to the interdigital space between the developing digits 1 and 2 and is adjacent to the expression of *SOX-9* in these two digits at stage 49 (Figure 1). At this stage, *SOX-9* expression correlates with the developing cartilage elements of the wrist (carpals) and the first two digits (phalanges) (Figure 1; Victoria blue of stage 49). At stage 50, *BMP-2* was mainly expressed between developing digits 2 and 3 and was adjacent to the expression of *SOX-9* in these two digits at stage 51 (Figure 1). At this stage, the expression of *SOX-9* at the tip of digit 2 correlated with the formation of skeletal elements in digit 2 (i.e. the phalanges) and the expression in digit 3 primordium was not yet associated with skeletal elements as revealed by Victoria blue staining (Figure 1; Victoria blue of stage 51). Finally, at stage 52, *BMP-2* was expressed between developing digits 3 and 4 and was adjacent to the expression of *SOX-9* in these two digits at stage 53 (Figure 1). At that stage, the expression of *SOX-9* at the tip of digit 3 correlated with the formation of skeletal elements in digit 3 but the expression in digit 4 primordium was not yet associated with skeletal elements as revealed by Victoria blue staining (Figure 1; Victoria blue of stage 53). In addition to its expression between digits 3 and 4 at stage 52, *BMP-2* expression was still detectable at the base of all 4 digits. This expression was not investigated further in the present study but has been associated with the elongation phase of digits in other species [53-55].

Effects of blocking *Shh* signaling in developing digits

As described above, at stage 47, *BMP-2* and *Shh* expression are on opposite sides of the developing limb bud (Figure 1; first column). It has been clearly demonstrated in mouse that *BMP-2* is not essential for digit identity [18]. It was shown (in knockout mice for *Shh*) that *Shh* is not essential for chondrogenesis and is dispensable for *BMP-2* expression [9]. Based on these observations, we tested, in axolotl, whether *BMP-2* expression is independent of *Shh* signaling. The hedgehog pathway antagonist cyclopamine [56] was used to determine whether *BMP-2* expression is independent of *Shh* signaling. The expression of *BMP-2* and *SOX-9* were assessed in limbs of animals treated with 2 different concentrations of cyclopamine that were shown in a previous study to prevent the formation of posterior digits [42]. In the aforementioned study, the digits that remained after cyclopamine treatment were identified as follows: when 2 digits remained after treatment, they




Figure 1 Expression patterns (whole mount *in situ* hybridisation) for *BMP-2*, *SOX-9* and *Shh* in developing limbs and cartilage staining (Victoria blue V.b.). All pictures show a dorsal view of the limb with the posterior side on top. Numbers in the upper left of each panel refers to the developmental stages [51].

were found to be digits 1 and 2, and when only 1 digit remained (the result of a higher concentration of cyclopamine), it was identified as digit 1 [42]. In the present study, cyclopamine treatment was initiated at developmental stage 41 which corresponds to the onset of *Shh* expression [40,51]. We confirmed that the expression of *Shh* at various limb development stages (data not shown) was as it had been reported previously by two other groups [39,40]. The group of Imokawa *et al.* also reported that *Shh* is expressed as late as the 2-3 digit stage [40] which corresponds to stage 51-52 based on Nye *et al.* [51]. The expression of *BMP-2* and *SOX-9* was assessed after 14 (stage 51) and 21 (stage 53) days of treatment to see whether their expression correlated with skeletogenesis even in the absence of *Shh* signaling. *BMP-2* expression was not affected by the presence of cyclopamine prior to stage 51 of limb development (see Additional File 2: Figure S2 for the expression of *BMP-2* at stage 45 in control and cyclopamine treated limbs). The expression of *SOX-9* was not affected either at stage 45 in cyclopamine treated animals (see Additional File 2: Figure S2).

14 days

After 14 days of treatment (Figure 2A-F), control animals (treated with ethanol, the carrier for cyclopamine) had reached stage 51 with digits 1 and 2 in formation

(Figure 2A, D). *BMP-2* was expressed in the interdigital space of digits 1-2 and 2-3 (Figure 2A) and *SOX-9* was expressed at the tips of digits 1 and 2, and fully expressed in the forming digit 3 and in the carpal elements of the wrist (Figure 2D). As expected, *BMP-2* expression was adjacent to *SOX-9* expression in digits 1, 2 and 3. In limbs treated with 1 $\mu\text{g}/\text{mL}$ of cyclopamine (Figure 2B, E), digits 1 and 2 were deformed and in animals treated with 2 $\mu\text{g}/\text{mL}$ of cyclopamine (Figure 2C, F), only 1 digit was formed. The expression pattern of *BMP-2* was adjacent to the expression of *SOX-9* in the forming digits of cyclopamine treated animals just as it is in control animals.

21 days

After 21 days of treatment (Figure 2G-O), control animals had reached stage 53, and at this stage the pattern is complete. Limbs treated with 1 $\mu\text{g}/\text{mL}$ of cyclopamine developed only digits 1 and 2 and limbs treated with 2 $\mu\text{g}/\text{mL}$ of cyclopamine developed only digit 1. In control animals, the expression of *BMP-2* was observed around the digits, and the expression of *SOX-9* was in digits 3 and 4. This developmental stage corresponds to the elongation phase of digits in other species [53-55] and as previously mentioned, was not investigated further. In the animals treated with 1 $\mu\text{g}/\text{mL}$ (Figure 2H, K, N) or 2 $\mu\text{g}/\text{mL}$ of cyclopamine (Figure 2I, L, O) the expression




Figure 2 Expression of *BMP-2* (A-C) and *SOX-9* (D-F) detected by whole mount *in situ* hybridisation in developing limbs treated 14 days with cyclopamine. Expression of *BMP-2* (G-I) and *SOX-9* (J-L) in developing limbs treated 21 days with cyclopamine. Left column (A, D, G, J, M) shows control animals (0.4 µL EtOH/mL), middle column (B, E, H, K, N) shows the animals treated with 1 µg/mL of cyclopamine and right column (C, F, I, L, O) shows the animals treated with 2 µg/mL of cyclopamine. All pictures show a dorsal view of the limbs with posterior side on top. Arrowheads in C, H and K indicate the expression of *BMP-2* on both sides of the regenerating digits and the expression of *SOX-9* in the middle of the regenerating digits. The last row (M-O) shows the cartilage staining (Victoria blue) of the skeletal elements at the end of cyclopamine treatments (21 days).

of *BMP-2* and *SOX-9* in the digits that have formed (digits 1 and 2 for 1 µg/mL treatment and digit 1 for 2 µg/mL treatment) was adjacent, with the expression of *BMP-2* surrounding the expression of *SOX-9* as in the digits of control animals.

***BMP-2* and *SOX-9* expression during limb regeneration**

Limb regeneration in salamander can be divided into two main phases. 1) A preparation phase that starts at

amputation time and focuses mainly on the formation of the blastema; and 2) a redevelopment phase that starts approximately at the late bud stage and focuses mainly on the differentiation and repatterning of the blastema into a new limb (see reference [57] for stages description). To determine when *BMP-2* and *SOX-9* are expressed, whole mount *in situ* hybridisation was performed for both genes throughout limb regeneration. The onset of *BMP-2* expression occurred at the early

bud stage (EB; Figure 3A) while *SOX-9* was not yet expressed (Figure 3D). The expression of *BMP-2* was maintained distally in the blastema of medium bud (MB) and late bud (LB) stages (Figure 3B-C) and was adjacent to the more proximal expression of *SOX-9*; which starts at MB stage (Figure 3E-F). At the Palette (Pal) stage, *BMP-2* (Figure 3G) and *SOX-9* (Figure 3J) expression started to localize in the anterior region where the first two digits eventually form (as revealed by Victoria blue staining; Figure 3N). At this stage, the expression of *BMP-2* (Figure 3G) versus that of *Shh* was on the opposite side of the autopod (Figure 3M). As in development at stage 47 (Figure 1), the opposing pattern of expression of *Shh* and *BMP-2* just preceded the onset of digit chondrogenesis. At Pal stage, *BMP-2* (Figure 3G) delineates the first 2 digits, and at early differentiation (ED) stage, *BMP-2* expression also appears between digits 3 and 4 (Figure 3H). At these two stages (Pal and ED), the expression of *SOX-9* (Figure 3J-K), was adjacent to *BMP-2* expression (Figure 3G-H) and corresponded to the exact location of the skeletal elements. In later stages (Figure 3K-L) *SOX-9* expression was tightly correlated to digit chondrogenesis as revealed by Victoria blue staining (Figure 3N-O).

Effects of blocking *Shh* signaling on regenerating digits

As described above, at Pal stage, *BMP-2* expression is on the opposite side of the regenerating limb bud compared to that of *Shh* (Figure 3G & 3M). To determine if *Shh* signaling had an effect on *BMP-2* expression during limb regeneration, as determined for axolotl developing limbs (Figure 2), axolotls were treated with cyclopamine. The expression of *BMP-2* and *SOX-9* (determined by RT-PCR at MB stage when *BMP-2*, *SOX-9* and *Shh* are first co-expressed) was not affected after 6 or 24 hours of exposure to 1 or 2 $\mu\text{g}/\text{mL}$ of cyclopamine (data not shown). The expression of *BMP-2* at medium bud was not affected in regenerating limbs exposed to cyclopamine from the time of amputation either (see Additional File 3: Figure S3). Therefore, in order to mimic as close as possible the *Shh* knockout phenotype, axolotls were treated from the moment of limb amputation until ED (when patterning is completed). The expression of *BMP-2* and *SOX-9* was assessed at ED stage to determine whether their expression correlated with skeletogenesis during the redevelopment phase of regeneration as shown for development (see Figure 2). In animals treated with 1 $\mu\text{g}/\text{mL}$ of cyclopamine, *BMP-2* expression (Figure 4B) was adjacent to *SOX-9* expression (Figure 4E). The same observation was made in the 2 $\mu\text{g}/\text{mL}$ treated animals (compare Figure 4C with 4F). Under these conditions, the expression of *BMP-2* and *SOX-9* relative to each other was the same as observed in corresponding digits of control limbs (Figure 4A & 4D). As in development, the 1 $\mu\text{g}/\text{mL}$

treated animals regenerated limbs with only two digits (Figure 4H) while the 2 $\mu\text{g}/\text{mL}$ treated animals regenerated limbs with only one digit (Figure 4I). Hence, the effect of cyclopamine on patterning regenerated skeletal elements as well as the expression of *BMP-2* and *SOX-9* were virtually identical to that observed in developing limbs treated with cyclopamine.

Overexpression of *BMP-2* and *Noggin* in regenerating limbs

Prior to investigating the *in vivo* role of *BMP-2* during limb regeneration, constructs expressing the monomeric red fluorescent protein (*mRFP*), *BMP-2* and *Noggin* were tested *in vitro* to verify their efficacy (Figure 5). C28/I2 cells were cotransfected with *mRFP*, axolotl *BMP-2* and *Xenopus laevis* *Noggin* along with the *BMP*-responsive BRE2-luc luciferase reporter construct [58] as well as a CMV- β -galactosidase construct as a control. Luciferase and β -galactosidase activities were determined 48 hours later and results (mean \pm sd; n = 3/condition) are presented as relative light unit (RLU) normalised with β -galactosidase. The first bar shows a basal activity (*mRFP* alone) of 4.6 ± 0.6 . The second bar shows that axolotl *BMP-2* activity is significantly ($p < 0.05$) higher (27.7 ± 3.4) than the basal activity (an increase of 6.0 fold). In the third bar, the activity of *Xenopus laevis* *Noggin* (general *BMP* inhibitor) alone (3.1 ± 0.3) is significantly ($p < 0.05$) lower than *BMP-2* activity and also significantly ($p < 0.05$) lower than basal activity. Finally, the fourth bar shows the activity (13.3 ± 1.9) resulting from *BMP-2* in the presence of *Noggin*. The activity of this combination (*BMP-2* + *Noggin*) is significantly ($p < 0.05$) lower than the *BMP-2* activity alone (a decrease of 2.0 fold). Comparison between each condition (n = 3/condition) was tested using Kruskal-Wallis One Way Analysis of Variance on Ranks. Hence, these results show the plasmids expressing axolotl *BMP-2* and *Xenopus laevis* *Noggin* produced functional proteins. As expected, the *BMP* responsive luciferase reporter construct was activated in the *BMP-2* transfected cells and was antagonized in the *Noggin* transfected cells (Figure 5).

In vivo experiments were performed on regenerating limbs and were based on the expression pattern of *BMP-2* & *SOX-9* determined by whole mount *in situ* hybridisation (Figure 3). *BMP-2* and *Noggin* containing plasmids were electroporated in the posterior-half of the blastema at the LB stages in order to obtain maximal expression of the transgene by the time the regenerates reached Pal; the stage when expression of *BMP-2* and *SOX-9* is localized in the anterior portion of the limb (Figure 3G & 3J). Hence, the electroporation was performed to induce *BMP-2* or *Noggin* at a time and place when no or low endogenous expression of *BMP-2* and *SOX-9* is detected.




Figure 3 Expression pattern, by whole mount *in situ* hybridisation, for *BMP-2* (A-C and G-I), *SOX-9* (D-F and J-L), *Shh* (M) and cartilage staining of skeletal elements (Victoria blue; N-O) at different stages of limb regeneration. All pictures show a dorsal view of the regenerating limbs with posterior side on the left. EB, early bud; MB, medium bud; LB, late bud; Pal, palette; ED, early differentiation; LD, late differentiation. The arrowhead in panel M indicates the restricted expression of *Shh* in the posterior portion of the blastema.





Figure 4 Expression pattern, by whole mount *in situ* hybridisation, for *BMP-2* (A-C) and *SOX-9* (D-F) at the ED stage in regenerating limb treated with cyclopamine from the time of amputation until fixed at ED. Left column (A, D, G) shows control animals (treated with the carrier 0.4 μL EtOH/mL), middle column (B, E, H) shows the animals treated with 1 μg/mL of cyclopamine and right column (C, F, I) shows the animals treated with 2 μg/mL of cyclopamine. All pictures show a dorsal view of the limbs with the posterior side on the left. The bottom row (G, H, I) show the skeletal staining (Victoria blue) at the end of the treatments when the control animals had completely regenerated.

The electroporation of *BMP-2* at the LB stage (Figure 6A) resulted in the absence of digit formation (Figure 6B-C) in the region of the electroporation ($n = 6/7$), while controls (electroporation of *mRFP* alone; Figure 6G) were normal (Figure 6H-I) in most cases ($n = 6/7$). In 1 control electroporated limb, 1 digit was missing as it sometimes happens in untreated animals (approximately 1% of the time, unpublished observations). The electroporation of *Xenopus laevis* *Noggin* under the same conditions (Figure 6D) resulted in a similar phenotype (Figure 6E-F; $n = 4/4$) to those obtained with the electroporation of *BMP-2*. When the same quantity of plasmid was injected throughout the blastema to obtain an even distribution of expression across the blastema with *BMP-2* ($n = 2/2$) or *Noggin* ($n = 3/4$), regeneration

was inhibited (data not shown). Expression of the *mRFP* construct alone throughout the blastema resulted in normally regenerating limbs ($n = 4/4$) therefore ruling out the electroporation itself as a cause for the inhibition of regeneration (data not shown). All electroporations performed prior to the LB stages with *BMP-2* in the full blastema had very mild effects ($n = 3/10$; digits were a little shorter in the electroporated region) and in the majority of cases, had no effect ($n = 7/10$) (data not shown). The electroporation with *Noggin* prior to the LB stage had no effect in all cases ($n = 12/12$) (data not shown). In control animals the electroporation of *mRFP* prior to LB gave rise to normal limbs ($n = 9/9$). With every construct combination used (*BMP-2/mRFP* or *Noggin/mRFP* or *mRFP* alone), the fluorescence detected



from the *mRFP* construct lasted a short time (5-7 days) when the electroporation was done prior to LB stage, compared to electroporation at LB stage which lasted longer (up to 15 days).

To investigate further the effects of *BMP-2* mis-expression, histological analysis was performed using the Masson's trichrome staining on limbs electroporated with *BMP-2* or *Noggin* (Figure 7A, D, G). Although the phenotypes of animals electroporated with *BMP-2* or *Noggin* appeared similar from an exterior perspective (Figure 6), the histology results showed significant differences at the tissue level. While limbs electroporated with *BMP-2* showed an increase in cellular density (possibly due to cellular condensation; n = 3) in the

electroporated region (Figure 7A), limbs electroporated with *Noggin* (n = 3) did not show any sign of such increase in cellular density or condensation (Figure 7D). Hence, these results suggest that the similarity between the phenotypes obtained with *BMP-2* and *Noggin* are caused by opposite effects of the ectopically expressed genes; an increase in cellular density or condensation with *BMP-2* and a lack of cellular density or condensation with *Noggin* leading to a loss of digits.

Histological analysis provided some indication as to why *BMP-2* and *Noggin* overexpression caused a loss of digits in regenerating limbs but it did not provide any mechanisms. Therefore, in order to help uncover the mechanism(s) as to how digit loss resulted following





Figure 6 Ectopic expression of axolotl *BMP-2* and *Xenopus laevis* *Noggin* by electroporating the expression plasmids directly into regenerating limbs. Different plasmid combinations were transfected to test the effect of *BMP-2* and *Noggin* on the regeneration process. In order to assess the efficiency of the electroporation in vivo (first column A, D, G), the *mRFP* expressing plasmid was co-transfected with axolotl *BMP-2* (A), *Xenopus laevis* *Noggin* (D) or *mRFP* alone (G; control) in the posterior half of regenerating blastema at the LB stage. The second column (B, E, H) show the resulting phenotypes once the controls (*mRFP* ectopic expression alone) were fully regenerated and the third column (C, F, I) show their skeletal elements using Victoria blue staining.

BMP-2 and *Noggin* overexpression, cell division and apoptosis were measured 6 days after electroporation of *mRFP*, *BMP-2* and *Noggin*. Electroporation was performed at LB stage as described above and cell division was determined by 5-bromo-2-deoxyuridine (BrdU) incorporation (n = 3) and apoptosis was determined by terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) assay (n = 3). *BMP-2* overexpression did not seem to cause major changes in cell division (Figure 7B) compared to the controls electroporated with *mRFP* (Figure 7H) which regenerated normally (Figure 7G). Although *BMP-2* overexpression did not

cause major changes in cell division, it did cause a marked increase in the number of apoptotic cells (Figure 7C) when compared to control electroporated with *mRFP* (Figure 7I). On the opposite, *Noggin* overexpression resulted in a marked reduction in cell division (Figure 7E) and the near absence of apoptotic cells (Figure 7F) compared to *BMP-2* overexpression and the control electroporated with *mRFP*.

Discussion

Our results show that the expression of *BMP-2* and *SOX-9* in developing axolotl digits is very similar to the



expression observed in other non amphibian species (e. g. mouse & chicken) [11,15,19,24,26,29,30,45-48]. The expression pattern of these two genes is also reproduced during digit formation in regenerating axolotl limbs. Since *SOX-9* is the earliest marker of chondrogenesis and a known target of *BMP-2*, these results suggest an activation of *SOX-9* expression (in cells that will form the digit bones) by *BMP-2* expressing cells (in the ID tissue).

As described by previous studies [42,43], the number of digits that form in cycloamine treated animals (during development and regeneration) is dose-dependent.

In the digits that do form in cycloamine treated animals, the expression of *BMP-2* and *SOX-9* as well as chondrogenesis proceed normally. These results suggest that in axolotl just as it was observed in mice [9], *BMP-2* is not likely to be a secondary signal for *Shh* signaling. This is also supported by the fact that when digits begin to condense in developing and regenerating limbs, *BMP-2* and *Shh* expression are on opposite sides of the limbs (along the anterior-posterior axis) and do not overlap. Hence, these results suggest that, as in mouse, *BMP-2* can trigger cell condensation independently of *Shh* signaling.

Our results strongly suggest that the inhibition of digit regeneration, in the limbs in which *BMP-2* was overexpressed, is the result of increased cell condensation and apoptosis. This is also supported by a previous study showing an inhibition of chick limb development in the presence of ectopic *BMP-2* [59]. The *BMP-2* overexpression experiments and the complementarities of expression of *BMP-2* and *SOX-9* (in developing and regenerating limbs), support our hypothesis that *BMP-2* triggers the condensation of adjacent cells destined to become chondrocytes (e.g. in the digits). Taken together, these results suggest that the role of *BMP-2* during limb regeneration is to trigger cell condensation and may be to control, to some extent, apoptosis. The data obtained with *Noggin* overexpression also provides further insight on the potential role of *BMP* signaling in the regeneration process. *Noggin* is an inhibitor of *BMPs* (general inhibitor of all *BMPs*) and therefore the results obtained could potentially affect *BMPs* other than *BMP-2*. However, these results indicate that *BMP* signaling is essential for cellular division and apoptosis during the regeneration process, since both processes are inhibited when *Noggin* is overexpressed.

The fact that the effects of *BMP-2* and *Noggin* were observed when overexpressed at the LB stage suggest that *BMP-2* regulates cellular activities during the redevelopment phase of regeneration that starts approximately at the LB stage or later (see [60,61] for a more complete description of the different phases of regeneration which are often referred to as: preparation phase which includes the initial stages leading to blastema formation; and redevelopment phase which includes the stages that give rise to the differentiated tissues and the patterning of the regenerated limb). However given that overexpression of *BMP-2* and *Noggin* were performed using electroporation of plasmids, technical limitations do not allow us to draw any conclusion regarding the role of *BMP-2* during the preparation phase of regeneration. Given the high level of cell proliferation and tissue reorganisation during the stages prior to the LB stage, it is possible that the electroporated plasmids could have been degraded before affecting the regeneration process or that cells compensated somehow to overcome the ectopic *BMP* signaling. This is supported by the rapid rate of fading of red fluorescence resulting from the reporter *mRFP* plasmid, when electroporations were done prior to LB stage. In addition, the fact that *Noggin* did not affect regeneration when overexpressed prior to LB stage is indicative that *BMP* signaling may not be essential for cellular proliferation during EB and MB stages.

Conclusion

This work represents the first study on the role of *BMPs* during the process of limb regeneration in urodele

amphibians. Our data demonstrate that the role of *BMP-2* is conserved in limb regeneration as in limb development and that its function is related to skeletogenesis, cellular proliferation and apoptosis but not pattern formation, as previously demonstrated by the group of Tabin in mice [18]. The ectopic expression of *BMP-2* led to the condensation of cells in regenerating axolotl limbs. The expression of *BMP-2* and *SOX-9* in addition to our functional analysis data from the ectopic expression of both *BMP-2* and *Noggin* clearly indicate that *BMP* signaling is essential for the process of limb regeneration especially during the redevelopment phase where re-differentiation of all the different cell types and pattern formation occur.

Methods

Animal maintenance and surgery

Axolotl embryos and larvae (3.0 cm to 8.5 cm from snout to tip of the tail) were purchased from the *Ambystoma* Genetic Stock Center (Lexington, KY, USA). Animals were kept in 20% Holtfreter's solution at a temperature varying from 19-22°C with a photoperiod cycle of 12 hours of light and 12 hours of darkness. Embryos were used for limb development experiments and larvae for limb regeneration experiments. For all manipulations performed, axolotls were anaesthetized in 0.1% MS222 (ethyl 3-aminobenzoate methanesulfonate salt, Sigma-Aldrich, MO, USA) in Holtfreter's solution at pH 7.0. Axolotls were allowed to recover from anaesthesia in 20% Holtfreter's or were euthanized/fixed for subsequent experiments. All animal care and experiments were done in agreement with the Université de Montréal animal care committee's guidelines.

Cloning of axolotl *BMP-2* full length cDNA

Partial axolotl *BMP-2* cDNA of 239 bp was obtained from axolotl larvae total RNA by RT-PCR. The 239 bp cDNA was amplified with primers (see Additional File 4: Figure S4) designed from a partial cDNA sequence of *BMP-2/4* like directly submitted to Genbank by Dalglish, G. *et al.* (Accession number: AY326272). The full length *BMP-2* cDNA was subsequently obtained by screening an axolotl cDNA library (Stratagene, CA, USA), following the manufacturer's instructions and using the 239 bp fragment as a probe radioactively labeled with ³²P-dCTP (Perkin Elmer, MA, USA).

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization was performed as described in Gardiner *et al.* [62] with a few modifications: The axolotls were anaesthetized and euthanized/fixed in 1× MEMFA (10× MEM salt (1 M MOPS pH 7.4, 20 mM EGTA, 10 mM MgSO₄), 3,7% formaldehyde in DEPC treated H₂O) then transferred to 100% methanol and

stored at -20°C until needed. Fragments of 449 bp and 381 bp (for *BMP-2* probes), 458 bp and 455 bp (for *SOX-9* probes) and 371 bp (for *Shh* probes) were generated by RT-PCR (see Additional File 4: Figure S4 for primers used) and cloned separately into the pCRII-TOPO cloning vector (Invitrogen, CA, USA). These plasmids were then linearized using the appropriate restriction enzymes and used as template for generating antisense RNA probes. Digoxigenin (DIG) labelled antisense RNA probes were generated using T7 or SP6 RNA polymerase (Invitrogen, CA, USA) and DIG RNA labelling mix (Roche Diagnostics, QC., Canada). Two different probes for *BMP-2* were synthesised. The 3' UTR of the *BMP-2* gene is highly conserved throughout evolution and it was utilized to make a highly specific probe for *BMP-2*. Thus, one probe of 381 bp was designed to overlap part of the 3' UTR and the ORF of *BMP-2*. A second probe of 449 bp was designed in the region overlapping the 5' UTR and the ORF of the gene which confirmed the expression observed with the 381 bp probe (data not shown). The expression pattern of *SOX-9* was also determined using two different probes with identical results obtained for both probes (data obtained using the 455 bp probe are not shown). The expression of *Shh* was assessed with only one probe since the results were identical to the results published by two different groups [39-41]. For tissue permeabilization of developing embryos, the limbs were incubated with 20 $\mu\text{g}/\text{mL}$ proteinase K for 15 min on ice and then at 37°C for 5 min. For permeabilization of regenerating tissues, limbs were incubated with 30 $\mu\text{g}/\text{mL}$ proteinase K for 1 h on ice and then at 37°C for 45 min. Probes were hybridized 24 h for embryos and 72 h for regenerating animals. Prehybridization and hybridization temperatures were done at: 55°C for the *BMP-2* 449 bp probe and 60°C for all the other probes. Finally, BM purple (Roche Diagnostics, QC., Canada) was used as the alkaline phosphatase substrate for the colorimetric reaction. Each time point presented were performed at least 3 times.

Whole mount skeletal staining of limbs

Cartilage was stained using the Victoria Blue method [63]. Limbs were fixed in Bouin's fixative solution for 24 h before being rinsed several times with 70% ethanol. Specimens were then rinsed in 3.5% NH_4OH for 2 days and subsequently treated with Acid alcohol for 2 h. Specimens were stained with 1% Victoria Blue for 2 h and then rinsed with 70% ethanol. Limbs were dehydrated using a gradient ethanol series of 95% and 100%, then cleared and stored in methyl salicylate.

Histology

Electroporated limbs were fixed overnight in Bouin's fixative solution and then rinsed thoroughly with 70%

alcohol. Limbs were embedded in paraffin and cut to 10 μm sections. Slides were deparaffinised through 3 baths of toluene for 5 minutes each. Slides were then rehydrated in a graded series of 100%, 90%, 70% and 50% ethanol and then distilled water for 5 minutes each. Masson's trichrome staining method was used to stain cell cytoplasm in red, nuclei in black and collagen in blue [64].

Cyclopamine treatments

Cyclopamine powder (Cedarlane, ON., Canada) was dissolved in 100% ethanol at a concentration of 5 mg/mL and stored at -20°C . Cyclopamine was administered by adding it directly to the Holtfreter's solution to the desired concentrations. For each conditions presented in this study, 10 animals were treated with cyclopamine or ethanol (control). For experiments on developing limbs, treatments began at the developmental stage 41 according to Nye *et al.* [51] and animals were fixed after 14 and 21 days of treatment (corresponding approximately to stage 51 and 53 in control animals). For regenerating animals, treatments began at the time of limb amputation and animals were fixed when the controls reached the palette stage (see [57] for stages description) and at completion of regeneration. For limb development as well as limb regenerating experiments, control animals were treated with the same volume of 100% ethanol that was used to dissolve the cyclopamine. Controls were done for both volumes of ethanol (0.2 $\mu\text{L}/\text{mL}$ and 0.4 $\mu\text{L}/\text{mL}$) used for the 1 and 2 $\mu\text{g}/\text{mL}$ cyclopamine treatment. There was no difference in limb development between these two control conditions (data not shown) and therefore only those corresponding to 2 $\mu\text{g}/\text{mL}$ cyclopamine treatment are presented. When the control animals with ethanol were compared with animals bathing in ethanol free solution, there was no difference in limb patterning and gene expression. All solutions were changed daily. Animals were euthanized at the end of the experiment and fixed.

Luciferase assays

The immortalized human chondrocyte cell line, C28/I2, has been described previously [65]. It was developed from chondrocytes isolated from juvenile human costal cartilage and immortalized by retroviral infection of the SV40 large T antigen [65]. C28/I2 cells were cultured in D-MEM/F12 containing 10% FBS with antibiotics (penstrep, fungizone) and 2 mM L-glutamine.

Transfection was performed using Lipofectamine 2000 (Invitrogen, Carlsbad, CA). BRE₂-luc was described previously [58]. C28/I2 cells were seeded at a density of 7.5×10^4 cells in a 12-well plate. The next day, cells were transfected with the following constructs and

concentrations: axolotl *BMP-2* = 0 or 0.4 μg , *Xenopus laevis* *Noggin* = 0 or 0.4 μg , $\text{BRE}_2\text{-luc}$ = 0.35 μg and pCMV- β gal = 0.35 μg . The total DNA concentration was brought up to 1.5 μg using pTYFP (expression vector for *mRFP*) as filler DNA. Cell lysates were prepared 48 hours later and analyzed for luciferase and β -galactosidase activity as described previously [66]. Data are expressed as a mean \pm standard deviation luciferase activity normalized to β -galactosidase activity. Comparison among each condition ($n = 3/\text{condition}$) was tested using Kruskal-Wallis One Way Analysis of Variance on Ranks.

Injections and electroporation of constructs

For injections/electroporation of constructs expressing axolotl *BMP-2* or *Xenopus laevis* *Noggin*, blastemas were injected with 1-1.5 μL of circular DNA solution through a glass needle with a 0.2-0.4 μm diameter tip, using a cell microinjector PM-1000 (Micro Data Instrument, Inc., NJ, USA). Immediately following injections, constructs were electroporated in blastemas. Thin gold plated rectangular electrodes (3 mm \times 5 mm) were placed under (anode) and over (cathode) the blastema. 5 trains (1 train/second) of 20 square wave pulses (200 Hz for 100 ms), at 10 Volts peak to peak, were applied through the electrodes using a S8 stimulator (Grass instruments, MA, USA; Additional File 5: Figure S5). Vectors containing axolotl *BMP-2* (5 $\mu\text{g}/\mu\text{L}$) or *Xenopus laevis* *Noggin* (5 $\mu\text{g}/\mu\text{L}$) were always co-injected with a vector containing the *mRFP* (2.8 $\mu\text{g}/\mu\text{L}$) to monitor the spread of the injection over time and space. Control animals were injected with 7.8 $\mu\text{g}/\mu\text{L}$ of *mRFP* (the vector containing *BMP-2* or *Noggin* was replaced by the same amount of *mRFP*). *BMP-2*, *Noggin* and *mRFP* expressions were all under the control of the constitutive CMV promoter in the pcDNA1/AMP expression vector (Invitrogen, CA).

BrdU incorporation assay

Regenerating axolotls were injected intra-peritoneally, 6 days after electroporation of plasmids (as described above); with 10 μL of BrdU stock solution using a ratio of 1-2 mL/100 g of body weight according to the manufacturer's instructions (GE Healthcare, # RPN201). Animals were fixed 12 hours after injection in 4% paraformaldehyde in 0.7 \times PBS for 24 hours at 4°C. Samples were paraffin embedded, deparaffinized and rehydrated as for histology. For immuno-histochemistry, slides were washed 4 \times 15 minutes in PBST (1 \times PBS with 0,1% Tween 20). They were incubated with 0.8% Pepsin in 0,2N HCl for 10 minutes at 37°C to promote denaturation of DNA. Slides were then washed 3 \times 10 minutes in PBST, and incubated in a blocking solution

(2% bovine serum albumin, 1% DMSO, 10% sheep serum and 0,1% Triton X-100) at room temperature for one hour. Slides were then incubated overnight at 4°C with an anti-BrdU mouse monoclonal antibody (BrdU Ab-3, Labvision/Neomarkers, Fremont, CA.) diluted 1:50 in blocking solution. PBST washes (4 \times 15 minutes) were done before incubating with an anti-mouse secondary antibody coupled to horseradish peroxidase, dilution 1:250, for 2 hours at room temperature (GE Healthcare). Slides were washed again 4 \times 15 minutes in PBST before being incubated in DAB (Zymed, Invitrogen CA) for signal detection. Slides were counterstained with methyl green for 6 minutes (Dako, Mississauga, ON) before being serially dehydrated 2 \times 2 minutes each in 90% EtOH, 100% EtOH and 100% Xylene. Slides were mounted with Permount (Fisher scientific, Ottawa, ON). Immunohistochemical detection for BrdU incorporation was done on three different samples for each condition tested.

TUNEL assay

TUNEL assay was adapted from Vlaskalin *et al.* [67] with minor modifications. Tissue sections were deparaffinized and rehydrated as for histology. In a humidified chamber, sections were incubated for 20 minutes in TBS (Tris-buffered saline: 100 mM Tris-HCL pH 7.5 + 150 mM NaCl), containing 20 $\mu\text{g}/\text{ml}$ proteinase K at RT and rinsed 3 \times 1 minute in TBS. Slides were rinsed in TBS, equilibrated in 1 \times TdT buffer for 5 minutes and incubated in TdT buffer containing 75 U/ml TdT and 2 μM digoxigenin-dUTP for 1 h at 37°C. End labelling was stopped by transferring the sections into 1 \times TBS/1 mM EDTA for 10 minutes at RT, followed by 2 \times 2 minutes rinses in TBS. Sections were equilibrated for 5 minutes in TBS and incubated in blocking solution [2% sheep serum in TBS] for 1 h at RT. Tissues were incubated overnight at 4°C in blocking solution containing 1:1500 dilution of anti-DIG-AP. The following day, sections were rinsed in PBS and equilibrated 5 minutes in AP buffer. The chromogenic reaction was carried out in AP buffer containing 100 mg/ml NBT and 50 mg/ml BCIP. The reaction was monitored for up to 15 minutes and stopped by several washes in 1 \times PBS.

Image treatments

Pictures of whole mounts, Victoria blue staining, and *mRFP* fluorescence were taken by a digital infinity 2 camera (Lumenera corporation, ON., Canada) through a Leica MZ-16F binocular (Leica Microsystems Ltd., Switzerland). Images of fluorescence in regenerating limbs were merged with white light images using Photoshop 7.0.1 (Adobe Systems Inc., CA, USA).

Additional file 1: Figure S1: Sequence analysis of axolotl BMP-2.

Sequence analysis and alignment of the axolotl *BMP-2* protein. Alignment of the predicted axolotl *BMP-2* protein with human, mouse and *Xenopus*. Seven cysteine residues that are highly conserved and characteristic to all members of the *Tgf-β* superfamily are shaded in red. Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-213X-10-15-S1.TIFF]

Additional file 2: Figure S2: Expression of BMP-2 and SOX-9 at stage 45 of limb development.

Expression of *BMP-2* and *SOX-9* at stage 45 of limb development in control and cyclopamine treated (2 μg/mL) axolotls. *Shh* has been shown to be expressed at this stage of axolotl limb development. Our results indicate that the expression of *BMP-2* and *SOX-9* during limb development is not dependent on *Shh* signaling as they are not affected by cyclopamine treatment. Panels A and B show the expression of *BMP-2* in control (A) and cyclopamine treated (B) animals. Panels C and D show the expression of *SOX-9* in control (C) and cyclopamine treated (D) animals. Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-213X-10-15-S2.TIFF]

Additional file 3: Figure S3: Expression of BMP-2 and SOX-9 at MB with cyclopamine.

Expression of *BMP-2* and *SOX-9* at MB stage of limb regeneration in control and cyclopamine treated (2 μg/mL) axolotls. *Shh* has been shown to be expressed at this stage of axolotl limb regeneration. Our results indicate that the expression of *BMP-2* during limb regeneration is not dependent on *Shh* signaling as it is not affected by cyclopamine treatment. Panels A and B show the expression of *BMP-2* in control (A) and cyclopamine treated (B) animals. Panels C and D show the expression of *SOX-9* in control (C) and cyclopamine treated (D) animals. Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-213X-10-15-S3.TIFF]

Additional file 4: Figure S4: Primers. Primers used for PCR amplification of the different probes used for whole mount in situ hybridization and cDNA library screening. Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-213X-10-15-S4.TIFF]

Additional file 5: Figure S5: In vivo electroporation. Schematic representation of the electric pulses used for electroporating the plasmids in regenerating tissues. 5 trains of pulses were applied for the *in vivo* electroporation of expression constructs. For each electroporation, a train of 20 square waves (10 V peak to peak) in 100 ms (200 Hz) is applied every second (1 Hz) over five seconds (5 trains in total). Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-213X-10-15-S5.TIFF]

Acknowledgements

The authors would like to thank Cara Hutchison and Mireille Pilote for the use of their plasmids containing the fragments for the two *Sox-9* probes used in this paper. We would also like to acknowledge the help of Éric Villiard and Samuel Gatien in preparing the photographs presented in the various figures. The authors would also like to thank Dr Antonio Nanci, from the department of Stomatology at the Université de Montréal, for help with the histology and Dr Marko Horb from the Institut de Recherche Clinique de Montréal, for providing the *Xenopus laevis* *Noggin* plasmid. This work was funded by a grant from the National Sciences and Engineering Research Council of Canada (NSERC) to SR. J-CG was supported by a Ph.D. scholarship from NSERC, P-LM and JB were supported by an undergraduate studentship from the Network for Oral Research Training and Health of the Canadian Institutes of Health Research and ML was supported by a Ph.D. scholarship from the Réseau de Recherche en Santé buccodentaire et osseuse of the Fond de la Recherche en Santé du Québec.

Author details

¹Department of Stomatology, Faculty of Dentistry, Université de Montréal, Montreal (QC) Canada. ²Department of Biochemistry, Faculty of Medicine, Université de Montréal, Montreal (QC) Canada. ³Department of Surgery, Faculty of Medicine, McGill University, Montreal (QC) Canada.

Authors' contributions

J-CG carried out the majority of manipulations, participated in the design of the study and drafted the manuscript. ML performed substantial work in the histology experiments and helped in reviewing the manuscript. P-LM cloned the axolotl *BMP-2* cDNA. JB managed the cyclopamine experiments on developing axolotls. KF performed the Luciferase assays experiments and helped in reviewing the manuscript. AP supervised the Luciferase experiments and helped in reviewing the manuscript. SR participated in the design of the present study, supervised all the experimentations and did all the major corrections of the final manuscript. All authors read and approved the final manuscript.

Received: 25 June 2009

Accepted: 12 February 2010 Published: 12 February 2010

References

1. Urist MR: Bone: formation by autoinduction. *Science* 1965, **150**:893-9.
2. Wozney JM: Overview of bone morphogenetic proteins. *Spine* 2002, **27**: S2-8.
3. Munoz-Sanjuan I, Brivanlou AH: Neural induction, the default model and embryonic stem cells. *Nat Rev Neurosci* 2002, **3**:271-80.
4. Suzuki A, Kaneko E, Ueno N, Hemmati-Brivanlou A: Regulation of epidermal induction by BMP2 and BMP7 signaling. *Dev Biol* 1997, **189**:112-22.
5. Riddle RD, Johnson RL, Laufer E, Tabin C: Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell* 1993, **75**:1401-16.
6. Honig LS, Summerbell D: Maps of strength of positional signalling activity in the developing chick wing bud. *J Embryol Exp Morphol* 1985, **87**:163-74.
7. Tickle C, Summerbell D, Wolpert L: Positional signalling and specification of digits in chick limb morphogenesis. *Nature* 1975, **254**:199-202.
8. Lopez-Martinez A, Chang DT, Chiang C, Porter JA, Ros MA, Simandl BK, Beachy PA, Fallon JF: Limb-patterning activity and restricted posterior localization of the amino-terminal product of Sonic hedgehog cleavage. *Curr Biol* 1995, **5**:791-6.
9. Litingtung Y, Dahn RD, Li Y, Fallon JF, Chiang C: Shh and Gli3 are dispensable for limb skeleton formation but regulate digit number and identity. *Nature* 2002, **418**:979-83.
10. McGlenn E, Tabin CJ: Mechanistic insight into how Shh patterns the vertebrate limb. *Curr Opin Genet Dev* 2006, **16**:426-32.
11. Geetha-Loganathan P, Nimmagadda S, Huang R, Scaal M, Christ B: Expression pattern of BMPs during chick limb development. *Anat Embryol (Berl)* 2006.
12. Laufer E, Nelson CE, Johnson RL, Morgan BA, Tabin C: Sonic hedgehog and Fgf-4 act through a signaling cascade and feedback loop to integrate growth and patterning of the developing limb bud. *Cell* 1994, **79**:993-1003.
13. Yang Y, Drossopoulou G, Chuang PT, Duprez D, Marti E, Bumcrot D, Vargesson N, Clarke J, Niswander L, McMahon A, et al: Relationship between dose, distance and time in Sonic Hedgehog-mediated regulation of anteroposterior polarity in the chick limb. *Development* 1997, **124**:4393-404.
14. Hogan BL: Bone morphogenetic proteins: multifunctional regulators of vertebrate development. *Genes Dev* 1996, **10**:1580-94.
15. Francis PH, Richardson MK, Brickell PM, Tickle C: Bone morphogenetic proteins and a signalling pathway that controls patterning in the developing chick limb. *Development* 1994, **120**:209-18.
16. Drossopoulou G, Lewis KE, Sanz-Ezquerro JJ, Nikbakht N, McMahon AP, Hofmann C, Tickle C: A model for anteroposterior patterning of the vertebrate limb based on sequential long- and short-range Shh signalling and Bmp signalling. *Development* 2000, **127**:1337-48.
17. Duprez DM, Kostakopoulou K, Francis-West PH, Tickle C, Brickell PM: Activation of Fgf-4 and HoxD gene expression by BMP-2 expressing cells in the developing chick limb. *Development* 1996, **122**:1821-8.

18. Bandyopadhyay A, Tsuji K, Cox K, Harfe BD, Rosen V, Tabin CJ: **Genetic Analysis of the Roles of BMP2, BMP4, and BMP7 in Limb Patterning and Skeletogenesis.** *PLoS Genet* 2006, **2**:e216.
19. Laufer E, Pizette S, Zou H, Orozco OE, Niswander L: **BMP expression in duck interdigital webbing: a reanalysis.** *Science* 1997, **278**:305.
20. Dahn RD, Fallon JF: **Interdigital regulation of digit identity and homeotic transformation by modulated BMP signaling.** *Science* 2000, **289**:438-41.
21. Ganan Y, Macias D, Dutarque-Coquillaud M, Ros MA, Hurlle JM: **Role of TGF beta s and BMPs as signals controlling the position of the digits and the areas of interdigital cell death in the developing chick limb autopod.** *Development* 1996, **122**:2349-57.
22. Chen Y, Zhao X: **Shaping limbs by apoptosis.** *J Exp Zool* 1998, **282**:691-702.
23. Zou H, Niswander L: **Requirement for BMP signaling in interdigital apoptosis and scale formation.** *Science* 1996, **272**:738-41.
24. Yokouchi Y, Sakiyama J, Kameda T, Iba H, Suzuki A, Ueno N, Kuroiwa A: **BMP-2/-4 mediate programmed cell death in chicken limb buds.** *Development* 1996, **122**:3725-34.
25. Macias D, Ganan Y, Sampath TK, Piedra ME, Ros MA, Hurlle JM: **Role of BMP-2 and OP-1 (BMP-7) in programmed cell death and skeletogenesis during chick limb development.** *Development* 1997, **124**:1109-17.
26. Weatherbee SD, Behringer RR, Rasweiler Jt, Niswander LA: **Interdigital webbing retention in bat wings illustrates genetic changes underlying amniote limb diversification.** *Proc Natl Acad Sci USA* 2006, **103**:15103-7.
27. Duprez D, Bell EJ, Richardson MK, Archer CW, Wolpert L, Brickell PM, Francis-West PH: **Overexpression of BMP-2 and BMP-4 alters the size and shape of developing skeletal elements in the chick limb.** *Mech Dev* 1996, **57**:145-57.
28. Pizette S, Niswander L: **Early steps in limb patterning and chondrogenesis.** *Novartis Found Symp* 2001, **232**:23-36.
29. Healy C, Uwanogho D, Sharpe PT: **Regulation and role of Sox9 in cartilage formation.** *Dev Dyn* 1999, **215**:69-78.
30. Pizette S, Niswander L: **BMPs are required at two steps of limb chondrogenesis: formation of prechondrogenic condensations and their differentiation into chondrocytes.** *Dev Biol* 2000, **219**:237-49.
31. Weston AD, Hoffman LM, Underhill TM: **Revisiting the role of retinoid signaling in skeletal development.** *Birth Defects Res C Embryo Today* 2003, **69**:156-73.
32. Bell DM, Leung KK, Wheatley SC, Ng LJ, Zhou S, Ling KW, Sham MH, Koopman P, Tam PP, Cheah KS: **SOX9 directly regulates the type-II collagen gene.** *Nat Genet* 1997, **16**:174-8.
33. Zehentner BK, Dony C, Burtcher H: **The transcription factor Sox9 is involved in BMP-2 signaling.** *J Bone Miner Res* 1999, **14**:1734-41.
34. Zhu J, Nakamura E, Nguyen MT, Bao X, Akiyama H, Mackem S: **Uncoupling Sonic hedgehog control of pattern and expansion of the developing limb bud.** *Dev Cell* 2008, **14**:624-32.
35. Vargas AO, Fallon JF: **The digits of the wing of birds are 1, 2, and 3. A review.** *J Exp Zool B Mol Dev Evol* 2005, **304**:206-19.
36. Shubin N, Alberch P: **A morphogenetic approach to the origin and basic organisation of the tetrapod limb.** *Evolutionary Biology* 1986, **20**:319-387.
37. Hinchliffe JR, Vorobyeva EI: **Developmental basis of limb homology in urodeles: heterochronic evidence from the primitive hynobiid family.** *Novartis Found Symp* 1999, **222**:95-105.
38. Franssen RA, Marks S, Wake D, Shubin N: **Limb chondrogenesis of the seepage salamander, *Desmognathus aeneus* (amphibia: plethodontidae).** *J Morphol* 2005, **265**:87-101.
39. Torok MA, Gardiner DM, Izpissua-Belmonte JC, Bryant SV: **Sonic hedgehog (shh) expression in developing and regenerating axolotl limbs.** *J Exp Zool* 1999, **284**:197-206.
40. Imokawa Y, Yoshizato K: **Expression of Sonic hedgehog gene in regenerating newt limb blastemas recapitulates that in developing limb buds.** *Proc Natl Acad Sci USA* 1997, **94**:9159-64.
41. Imokawa Y, Yoshizato K: **Expression of sonic hedgehog gene in regenerating newt limbs.** *Wound Repair Regen* 1998, **6**:366-70.
42. Stopper GF, Wagner GP: **Inhibition of Sonic hedgehog signaling leads to posterior digit loss in *Ambystoma mexicanum*: parallels to natural digit reduction in urodeles.** *Dev Dyn* 2007, **236**:321-31.
43. Roy S, Gardiner DM: **Cyclopamine induces digit loss in regenerating axolotl limbs.** *J Exp Zool* 2002, **293**:186-90.
44. Roy S, Gardiner DM, Bryant SV: **Vaccinia as a tool for functional analysis in regenerating limbs: ectopic expression of Shh.** *Dev Biol* 2000, **218**:199-205.
45. Pajni-Underwood S, Wilson CP, Elder C, Mishina Y, Lewandoski M: **BMP signals control limb bud interdigital programmed cell death by regulating FGF signaling.** *Development* 2007, **134**:2359-68.
46. Beck CW, Christen B, Barker D, Slack JM: **Temporal requirement for bone morphogenetic proteins in regeneration of the tail and limb of *Xenopus tadpoles*.** *Mech Dev* 2006, **123**:674-88.
47. Akiyama H, Chaboissier MC, Martin JF, Schedl A, de Crombrughe B: **The transcription factor Sox9 has essential roles in successive steps of the chondrocyte differentiation pathway and is required for expression of Sox5 and Sox6.** *Genes Dev* 2002, **16**:2813-28.
48. Akiyama H, Stadler HS, Martin JF, Ishii TM, Beachy PA, Nakamura T, de Crombrughe B: **Misexpression of Sox9 in mouse limb bud mesenchyme induces polydactyly and rescues hypodactyly mice.** *Matrix Biol* 2007, **26**:224-33.
49. Gomez-Escobar N, Lewis E, Maizels RM: **A novel member of the transforming growth factor-beta (TGF-beta) superfamily from the filarial nematodes *Brugia malayi* and *B. pahangi*.** *Exp Parasitol* 1998, **88**:200-9.
50. Massagué J: **The transforming growth factor-beta family.** *Annu Rev Cell Biol* 1990, **6**:597-641.
51. Nye HL, Cameron JA, Chernoff EA, Stocum DL: **Extending the table of stages of normal development of the axolotl: limb development.** *Dev Dyn* 2003, **226**:555-60.
52. Christen B, Slack JM: **All limbs are not the same.** *Nature* 1998, **395**:230-1.
53. De Luca F, Barnes KM, Uyeda JA, De-Levi S, Abad V, Palese T, Mericq V, Baron J: **Regulation of growth plate chondrogenesis by bone morphogenetic protein-2.** *Endocrinology* 2001, **142**:430-6.
54. Minina E, Wenzel HM, Kreschel C, Karp S, Gaffield W, McMahon AP, Vortkamp A: **BMP and *Ihh*/PTHrP signaling interact to coordinate chondrocyte proliferation and differentiation.** *Development* 2001, **128**:4523-34.
55. Sears KE, Behringer RR, Rasweiler Jt, Niswander LA: **Development of bat flight: morphologic and molecular evolution of bat wing digits.** *Proc Natl Acad Sci USA* 2006, **103**:6581-6.
56. Chen JK, Taipale J, Cooper MK, Beachy PA: **Inhibition of Hedgehog signaling by direct binding of cyclopamine to Smoothened.** *Genes Dev* 2002, **16**:2743-8.
57. Iten LE, Bryant SV: **Forelimb regeneration from different levels of amputation in the newt, *Notophthalmus viridescens*: length, rate, and stages.** *Wilhelm Roux' archiv* 1973, **173**:p263-282.
58. Korczynski O, ten Dijke P: **Identification and functional characterization of distinct critically important bone morphogenetic protein-specific response elements in the *Id1* promoter.** *J Biol Chem* 2002, **277**:4883-91.
59. Niswander L, Martin GR: **FGF-4 and BMP-2 have opposite effects on limb growth.** *Nature* 1993, **361**:68-71.
60. Roy S, Levesque M: **Limb regeneration in axolotl: is it superhealing?.** *ScientificWorldJournal* 2006, **6**(Suppl 1):12-25.
61. Gardiner DM, Carlson MR, Roy S: **Towards a functional analysis of limb regeneration.** *Semin Cell Dev Biol* 1999, **10**:385-93.
62. Gardiner DM, Blumberg B, Komine Y, Bryant SV: **Regulation of HoxA expression in developing and regenerating axolotl limbs.** *Development* 1995, **121**:1731-41.
63. Bryant SV, Iten LE: **The regulative ability of the limb regeneration blastema of *Notophthalmus viridescens*: Experiments in situ.** *Wilhelm Roux's Archives of Developmental Biology* 1974, **174**:90-101.
64. Luna L: **Manual of histologic staining methods of the Armed Forces Institute of Pathology.** New York: Blakiston Division (McGraw Hill), Third 1968.
65. Goldring MB, Birkhead JR, Suen LF, Yamin R, Mizuno S, Glowacki J, Arbisser JL, Apperley JF: **Interleukin-1 beta-modulated gene expression in immortalized human chondrocytes.** *J Clin Invest* 1994, **94**:2307-16.
66. Finsson KW, Parker WL, ten Dijke P, Thorikay M, Philip A: **ALK1 opposes ALK5/Smad3 signaling and expression of extracellular matrix components in human chondrocytes.** *J Bone Miner Res* 2008, **23**:896-906.
67. Vlaskalin T, Wong CJ, Tsilifidis C: **Growth and apoptosis during larval forelimb development and adult forelimb regeneration in the newt (*Notophthalmus viridescens*).** *Dev Genes Evol* 2004, **214**:423-31.

doi:10.1186/1471-213X-10-15

Cite this article as: Guimond et al.: BMP-2 functions independently of SHH signaling and triggers cell condensation and apoptosis in regenerating axolotl limbs. *BMC Developmental Biology* 2010 **10**:15.